

Rehabilitation s/p Surgical Meniscal Repair

Saint Louis University – SSM Health Physical Therapy Orthopedic Residency
in Collaboration with SLUCare Physicians



Physician Referral for Physical Therapy for:

Patient Name: _____ **Date:** _____

Referring DX: Meniscal Tear (medial / lateral)

Recommended Frequency: 1 – 3 visits/ wk

Total Duration: 4 months

These guidelines, treatments, and milestones have been established to assist in guiding rehabilitation based on the most current available evidence. They are not intended to be substitute for sound clinical judgement with consideration of the individual contextual features of the patient and the demands of various functions/sports.

Recommendations	Precautions:*	The following factors may affect prognosis
<ul style="list-style-type: none"> • When implementing the below guidelines for rehabilitation of meniscal repairs with concomitant procedures, consider the following: <ul style="list-style-type: none"> ○ With ACL reconstruction: <ul style="list-style-type: none"> ✓ Promote protection of the ACL graft by limiting excessive anterior tibial translation ○ With ACL and MCL repair: <ul style="list-style-type: none"> ✓ Limit excessive anterior tibial translation and avoid valgus stress ○ With PCL reconstruction: <ul style="list-style-type: none"> ✓ Avoid aggressive posterior tibial translation • Use of the Soreness Rules⁶ when determining exercise progression 	<ul style="list-style-type: none"> • No loaded knee flexion beyond 45° until week 5¹⁻³ • No loaded knee flexion beyond 90° until week 8 • No forced knee hyperextension if anterior horn repair • No forced knee flexion if posterior horn repair • Avoid OKC exercise from 0-30° and CKC exercise from 90-120° if patient shows signs/symptoms of patellofemoral irritation^{4,5} 	<ul style="list-style-type: none"> • Shorter meniscus healing time if concomitant cruciate repair^{7,8} • Biopsychosocial factors such as pain catastrophizing, fear-avoidance behavior, and exercise self-efficacy



Rehabilitation s/p Surgical Meniscal Repair

Timeline	Milestones	Treatment Recommendations
<u>Week 1-2</u> (Day 0-14)	Active full knee extension AROM flexion to 90°	Amb WBAT knee brace locked 0° Supervised loaded flexion between 0-45° Core stabilization Hip strengthening Patellar mobilizations NMES as needed ⁹
<u>Weeks 3-4</u> (Day 15-28)	AROM 0-120° Full scar mobility Patellar mobility WNL Zero to trace effusion (Stroke Test ¹⁰)	Amb WBAT knee brace locked 0° Gait training Alter-G Treadmill Core stabilization Hip strengthening Stair progression
<u>Weeks 5-7</u> (Day 29-49)	AROM to WNL Normal gait No Effusion ≤ 2 errors on SL Squat ^{11,12} 5xSTS ≤ 1 SD of norms ¹³	WBAT Loaded flexion between 0-90° Gait training SL motor control CKC Core stabilization Hip strengthening
<u>Weeks 8-11</u> (Day 50-77)	≤ 1 errors on SL Squat Mod SEBT symmetry ≤ 4cm ¹⁴	Loaded flexion > 90° Running progression Strength and conditioning CKC Core stabilization Hip strengthening
<u>Weeks 12- Return-to-sport</u>	Hop tests symmetry > 90% ¹⁵ Zero errors on SL Squat Acute-to-chronic workload ratio < 1.5 ¹⁶⁻¹⁸	Functional hop tests Sport-specific drills Agility drills CKC Core stabilization Hip strengthening

Abbreviations: SL = single limb; CKC = closed kinetic chain; SD = standard deviation; Mod SEBT = modified Star Excursion Balance Test

<p>For questions regarding the patient’s medical care, new orders, or insurance questions: please contact your physician's office directly</p>	<p>For additional questions, comments, or concerns regarding the implementation of these physical therapy guidelines, please contact Chris Sebelki, PT, DPT, PhD, OCS, Director of the SLU – SSM Health Physical Therapy Residency 314 977 8724 OR chris.sebelki@health.slu.edu</p>
---	--

Rehabilitation s/p Surgical Meniscal Repair

Tests/Measures:

- Soreness Rules⁶

Criterion	Action
1. Soreness during warm-up that continues	2 days off, drop down 1 step
2. Soreness during warm-up that goes away	Stay at step that led to soreness
3. Soreness during warm-up that goes away and redevelops during session	2 days off, drop down 1 step
4. Soreness the day after lifting (not muscle soreness)	1 day off, do not advance program to the next step
5. No soreness	Advance 1 step per week or as instructed by healthcare professional

- Single Leg Squat^{11,12}

Movement Impairment	
Midfoot collapse	Early heel rise
Femoral adduction, IR	Pelvic drop
Poor control of knee when rising up	Excessive trunk flexion or knee extension on rising up

* Table adapted from Liebenson 2002 in Bailey et al 2010

- 5xSTS Normative Values¹³

Age (n)	Mean ± SD (95%CI)	Min-Max
14–19 (25)	6.5 ± 1.2 (6.0–7.0)	4.7–9.7
20–29 (36)	6.0 ± 1.4 (5.6–6.5)	3.9–11.2
30–39 (22)	6.1 ± 1.4 (5.5–6.8)	4.1–10.4
40–49 (15)	7.6 ± 1.8 (6.6–8.6)	5.6–13.2
50–59 (20)	7.7 ± 2.6 (6.5–8.9)	4.2–12.1
60–69 (25)	7.8 ± 2.4 (6.8–8.7)	4.7–15.1
70–79 (24)	9.3 ± 2.1 (8.4–10.1)	5.5–13.3
80–85 (14)	10.8 ± 2.6 (9.3–12.3)	5.8–17.6

- Return to sport dosing should consider Acute-to-chronic workload¹⁶⁻¹⁸
 - Each session calculated by multiplying RPE (0-10) by duration (minutes) to obtain workload (augmented units). For example, *RPE of 6 x 60 minutes = workload of 360 AUs.*
 - Acute workload = average workload over the course of 1 week
 - Chronic workload = average workload over course of 4 weeks

Rehabilitation s/p Surgical Meniscal Repair

References:

1. Kawai Y, Fukubayashi T, Nishino J. Meniscal suture. An experimental study in the dog. *Clin Orthop Relat Res* 1989;243:286-292.
2. Dowdy P et al. The effect of cast immobilization on meniscal healing. An experimental study in the dog. *Am J Sports Med* 1995; 23:721-728.
3. Zhongan Z et al. Treatment of longitudinal injuries in avascular area of meniscus in dogs by rephination. *Arthroscopy* 1988; 4:151-156.
4. Steinkamp LA et al. Biomechanical considerations in patellofemoral joint rehabilitation. *Amer J Sports Med.* 1993;23(3):438-444.
5. Escamilla RF. Knee biomechanics of the dynamic squat exercise. *Med Sci Sport Exer.* 2001;33(1):127-141.
6. Fees M, Decker T, Snyder-Mackler L, Axe MJ. Upper extremity weight-training modifications for the injured athlete. A clinical perspective. *Am J Sports Med.* 1998;26:732-742.
7. Barber A, Click S. Meniscus repair rehabilitation with concurrent anterior cruciate reconstruction. *Arthroscopy.* 1997;13(4):433-437.
8. Noyes FR, Barber-Westin SD. Arthroscopic repair of meniscus tears extending into the avascular zone with or without anterior cruciate ligament reconstruction in patients 40 years of age and older. *Arthroscopy.* 2000;16:822-829.
9. Kim K, Croy T, Hertel J, Saliba S. Effects of neuromuscular electrical stimulation after anterior cruciate ligament reconstruction on quadriceps strength, function, and patient-oriented outcomes: a systematic review. *J Ortho Sports Phys Ther.* 2010;40(7):383-391.
10. Sturgill, L., Snyder-Mackler, L., Manal, T., & Axe, M. (2009). Interrater reliability of a clinical scale to assess knee joint effusion. *J Ortho Sports Phys Ther.* 2009;39(12):845-849.
11. Liebenson C. Functional exercises. *J Bodywork and Movem Ther.* 2002;6(2):108-13
12. Bailey R, Selfe, J, Richards J. The single leg squat test in the assessment of musculoskeletal function: a review. *Physiother Ireland.* 2010;13(1):18-23.
13. Bohannon RW et al. Sit-to-stand test: performance and determinants across the age-span. *Isokinet Exerc Sci.* 2010;18(4);235-240.
14. Garrison J, Bothwell J, Wolf G, et al. Y balance test anterior reach symmetry at three months is related to single leg functional performance at time of return to sports following anterior cruciate ligament reconstruction. *Int J Sports Phys Ther.* 2015;(10)5:602-611.
15. Fitzgerald GK, Lephart SM, Hwang JH, Wainner RS. Hop tests as predictors of dynamic knee stability. *J Orthop Sports Phys Ther.* 2001;31:588-597.
16. Hulin BT, Gabbett TJ, Blanch P, et al. Spikes in acute workload are associated with increased injury risk in elite cricket fast bowlers. *Br J Sports Med.* 2014;48:708-712.
17. Hulin BT, Gabbett TJ, Caputi P, et al. Low chronic workload and the acute:chronic workload ratio are more predictive of injury than between-match recovery time: a two-season prospective cohort study in elite rugby league players. *Br J Sports Med.* 2016;50:1008–12.
18. Gabbett TJ, Kennelly S, Sheehan J, et al. If overuse injury is a ‘training load error’, should undertraining be view the same way? *Br J Sports Med.* 2016;50:1017-1018.